| Grade 7 | Grade 7 Yearlong Mathematics Map | | | | | | | |
|---|--|--|-------------------------------|---|--|--|--|--|
| Resources: | Resources: Approved from Board of Education Assessments: PARCC Assessments, Performance Series, District Benchmark Assessments | | | | | | | |
| 1. Make sense of problems and persevere in solv | | | | oning. | | | | |
| Domain | Cluster | Common Core Standard | Content | Skills | Vocabulary | | | |
| RP | Analyze proportional relationships and use them to solve real- world and mathematical problems. | | Ratio Unit Rate | 7.RP.1 Compute unit rates with ratio of fractions including ratios of length, areas, and other quantities of like or different units. | Complex fraction Ratio Unit Rate | | | |
| RP | Analyze proportional relationships and use them to solve real- world and mathematical problems. | 7.RP.2 Recognize and represent proportional relationships between quantities. | Proportional Relationships | 7.RP.2 | Proportion (equivalent ratio) Proportional relationship Scale factor | | | |
| RP | Analyze proportional relationships and use them to solve real- world and mathematical problems. | 7.RP.2a Decide whether two quantities are in a proportional relationship, e.g., by testing for equivalent ratios in a table or graphing on a coordinate plane and observing whether the graph is a straight line through the origin. | Proportional Relationships | 7.RP.2a Decide whether two quantities are in a proportional relationship by using a table or graphing on a coordinate plane. | Proportional Relationship Proportion Coordinate Plane Origin x-coordinate y-coordinate quadrant x-axis y-axis scale | | | |

| Domain | Cluster | Common Core Standard | Content | Skills | Academic Vocabulary |
|--------|--|--|-------------------------------|--|--|
| RP | Analyze proportional relationships and use them to solve real- world and mathematical problems. | 7.RP.2b Identify the constant of proportionality (unit rate) in tables, graphs, equations, diagrams, and verbal descriptions of proportional relationships. | Unit Rate | 7.RP.2b Derive the constant of proportionality (unit rate) in tables, graphs, equations, diagrams, and verbal descriptions of proportional relationships. | Unit Rate Table Graph Diagrams Equations |
| RP | Analyze proportional relationships and use them to solve real- world and mathematical problems. | | Proportional Relationships | 7.RP.2c Write equations to represent proportional relationships. | Proportional relationships Equation Proportion |
| RP | Analyze proportional relationships and use them to solve real- world and mathematical problems. | 7.RP.2d Explain what a point (x, y) on the graph of a proportional relationship means in terms of the situation, with special attention to the points (0, 0) and (1, r) where r is the unit rate. | - | 7.RP.2d Explain what a point (x, y) on the graph of a proportional relationship means in terms of the situation, with special attention to the points (0, 0) and (1, r) where r is the unit rate. | Unit Rate Rate of change (slope) Graphs Equations Origin Ordered pair x-axis y-axis x coordinate y coordinate Coordinate plane |

| Domain | Cluster | Common Core Standard | Content | Skills | Academic Vocabulary |
|--------|----------------------------------|---|-------------------------------|---|--|
| RP | them to solve real- world and | 7.RP.3 Use proportional relationships to solve multistep ratio and percent problems. Examples: simple interest, tax, markups and markdowns, gratuities and commissions, fees, percent increase and decrease, percent error. | Proportional Relationships | using proportional relationships. | Proportional relationship Proportion Ratio Percent Percent error Simple Interest Percent increase Percent decrease Markup Sales Tax Commissions Gratutidies |
| NS | previous understandings of | 7.NS.1 Apply and extend previous understandings of addition and subtraction to add and subtract rational numbers; represent addition and subtraction on a horizontal or vertical number line diagram. | Rational Numbers | 7.NS.1 | |
| NS | previous understandings of | 7.NS.1a Describe situations in which opposite quantities combine to make 0. For example, a hydrogen atom has 0 charge because its two constituents are oppositely charged. | Rational Numbers | 7.NS.1a Describe situations in which opposite quantities combine to make 0. | Property of Opposites |

| Domain | Cluster | Common Core Standard | Content | Skills | Academic Vocabulary |
|--------|---|--|---------------------------------|---|---|
| NS | Apply and extend previous understandings of operations with fractions to add, subtract, multiply, and divide rational numbers. | 7.NS.1b Understand p + q as the number located a distance q from p, in the positive or negative direction depending on whether q is positive or negative. Show that a number and its opposite have a sum of 0 (are additive inverses). Interpret sums of rational numbers by describing real-world contexts. | Rational Numbers | | Rational numbers Vertical number line Horizontal number line Property of Opposites Positive number Negative number Combine Number line |
| NS | | 7.NS.1b Understand p + q as the number located a distance q from p, in the positive or negative direction depending on whether q is positive or negative. Show that a number and its opposite have a sum of 0 (are additive inverses). Interpret sums of rational numbers by describing real-world contexts. | Rational Numbers | 7.NS.1b Prove that a number and its opposite have a sum of 0 (are additive inverses). | Rational number Absolute value Combine (Additive Inverses) Property of opposites |
| NS | | 7.NS.1b Understand p + q as the number located a distance q from p, in the positive or negative direction depending on whether q is positive or negative. Show that a number and its opposite have a sum of 0 (are additive inverses). Interpret sums of rational numbers by describing real-world contexts. | Rational Numbers | 7.NS.1b Interpret sums of rational numbers by describing real-world contexts. | Rational number |
| NS | Apply and extend previous understandings of operations with fractions to add, subtract, multiply, and divide rational numbers. | 7.NS.1c Understand subtraction of rational numbers as adding the additive inverse, $p - q = p + (-q)$. Show that the distance between two rational numbers on the number line is the absolute value of their difference, and apply this principle in real-world contexts. | Subtraction Rational numbers | 7.NS.1c Explain subtraction of rational numbers as adding the additive inverse | Rational number Combine (Additive Inverses) Absolute Value |

| Domain | Cluster | Common Core Standard | Content | Skills | Academic Vocabulary |
|--------|---|---|---|--|---|
| NS | understandings of operations with | 7.NS.1c Understand subtraction of rational numbers as adding the additive inverse, $p - q = p + (-q)$. Show that the distance between two rational numbers on the number line is the absolute value of their difference, and apply this principle in real-world contexts. | Subtraction Rational numbers | 7.NS.1c Prove that the distance between two rational numbers on the number line is the absolute value of their difference. | Rational number Absolute Value Number line Positive numbers Negative numbers |
| NS | Apply and extend previous understandings of operations with fractions to add, subtract, multiply, and divide rational numbers. | 7.NS.1d Apply properties of operations as strategies to add and subtract rational numbers. | Properties of Operations | 7.NS.1d Apply properties of operations as strategies to add and subtract rational numbers. | Rational number Associative Property Commutative Property Additive Identity Property of Opposites |
| NS | previous | 7.NS.2 Apply and extend previous understandings of multiplication and division and of fractions to multiply and divide rational numbers. | Multiplication and Division of Rational Numbers | | |
| NS | previous understandings of operations with fractions to add, subtract, multiply, and | 7.NS.2a Understand that multiplication is extended from fractions to rational numbers by requiring that operations continue to satisfy the properties of operations, particularly the distributive property, leading to products such as $(-1)(-1) = 1$ and the rules for multiplying signed numbers. Interpret products of rational numbers by describing real-world contexts. | | 7.NS.2a Apply the properties of multiplication and rules for multiplying signed numbers to rational numbers. | Rational number Distributive Property Associative Property Commutative Property Multiplicative Identity Fraction Signed numbers |

| Domain | Cluster | Common Core Standard | Content | Skills | Academic Vocabulary |
|--------|---|---|------------------------------------|---|--|
| NS | Apply and extend previous understandings of operations with fractions to add, subtract, multiply, and divide rational numbers. | 7.NS.2a Understand that multiplication is extended from fractions to rational numbers by requiring that operations continue to satisfy the properties of operations, particularly the distributive property, leading to products such as $(-1)(-1) = 1$ and the rules for multiplying signed numbers. Interpret products of rational numbers by describing real-world contexts. | | 7.NS.2a Interpret products of rational numbers by describing real-world contexts. | Rational number Product |
| NS | Apply and extend previous understandings of operations with fractions to add, subtract, multiply, and divide rational numbers. | 7.NS.2b Understand that integers can be divided, provided that the divisor is not zero, and every quotient of integers (with non-zero divisor) is a rational number. If p and q are integers, then $-(p/q) = (-p)/q = p/(-q)$. Interpret quotients of rational numbers by describing real- world contexts. | Divide rational numbers | 7.NS.2b Explain that integers can be divided, provided that the divisor is not zero, and every quotient of integers (with non-zero divisor) is a rational number. | Integers Rational numbers divisors quotients Undefined quotient |
| NS | Apply and extend previous understandings of operations with | 7.NS.2b Understand that integers can be divided, provided that the divisor is not zero, and every quotient of integers (with non-zero divisor) is a rational number. If p and q are integers, then $-(p/q) = (-p)/q = p/(-q)$. Interpret quotients of rational numbers by describing real- world contexts. | Divide rational numbers | 7.NS.2b Interpret quotients of rational numbers by describing real-world contexts. | Quotient Rational numbers |
| NS | Apply and extend previous understandings of operations with fractions to add, subtract, multiply, and divide rational numbers. | 7.NS.2c Apply properties of operations as strategies to multiply and divide rational numbers. | Multiplication Rational Numbers | 7.NS.2c Apply properties of operations as strategies to multiply rational numbers. | Rational number Associative Property Commutative Property Multiplicative Identity |

| Domain | Cluster | Common Core Standard | Content | Skills | Academic Vocabulary |
|--------|---|---|----------------------------|--|---|
| NS | Apply and extend previous understandings of operations with fractions to add, subtract, multiply, and divide rational numbers. | 7.NS.2c Apply properties of operations as strategies to multiply and divide rational numbers. | divide rational numbers | 7.NS.2c Apply properties of operations as strategies to divide rational numbers. | Rational number Associative Property Commutative Property Multiplicative Identity |
| NS | previous | 7.NS.2d Convert a rational number to a decimal using long division; know that the decimal form of a rational number terminates in 0s or eventually repeats. | rational numbers | 7.NS.2d Convert a rational number to a decimal using long division | Rational number decimal place value terminating decimal repeating decimal bar notation |
| NS | previous | 7.NS.2d Convert a rational number to a decimal using long division; know that the decimal form of a rational number terminates in 0s or eventually repeats. | rational numbers | 7.NS.2d Recognize that the decimal form of a rational number terminates in 0s or eventually repeats. | Rational number decimal place value terminating decimal repeating decimal bar notation |
| NS | Apply and extend previous understandings of operations with fractions to add, subtract, multiply, and divide rational numbers. | 7.NS.3 Solve real-world and mathematical problems involving the four operations with rational numbers. | rational numbers | 7.NS.3 Solve real-world and mathematical problems involving the four operations with rational numbers including complex fractions. | Rational number Complex fractions |

| Domain | Cluster | Common Core Standard | Content | Skills | Academic Vocabulary |
|--------|---|--|---------------------------|--|---|
| EE | Use properties of operations to generate equivalent expressions. | 7.EE.1 Apply properties of operations as strategies to add, subtract, factor, and expand linear expressions with rational coefficients. | Equivalent Expressions | rational coefficients, using properties of operations. | Associative Property Commutative Property Distributive Property Linear expression Rational coefficent |
| EE | Use properties of operations to generate equivalent expressions. | 7.EE.1 Apply properties of operations as strategies to add, subtract, factor, and expand linear expressions with rational coefficients. | Equivalent Expressions | rational coefficients, using properties of operations. | Associative Property Commutative Property Distributive Property Linear expression Factor Rational coefficent |
| EE | Use properties of operations to generate equivalent expressions. | 7.EE.2 Understand that rewriting an expression in different forms in a problem context can shed light on the problem and how the quantities in it are related. For example, a + 0.05a = 1.05a means that "increase by 5%" is the same as "multiply by 1.05." | Equivalent Expressions | problem context to shed light on the problem and how the quantities in it are related. | Equivalent ratio expression equation decimal percent fraction |

| Domain | Cluster | Common Core Standard | Content | Skills | Academic Vocabulary |
|--------|---|---|---|--|---|
| EE | Solve real-life and mathematical problems using numerical and algebraic expressions and equations. | 7.EE.3 Solve multi-step real-life and mathematical problems posed with positive and negative rational numbers in any form (whole numbers, fractions, and decimals), using tools strategically. Apply properties of operations to calculate with numbers in any form; convert between forms as appropriate; and assess the reasonableness of answers using mental computation and estimation strategies. For example: If a woman making \$25 an hour gets a 10% raise, she will make an additional 1/10 of her salary an hour, or \$2.50, for a new salary of \$27.50. If you want to place a towel bar 9 3/4 inches long in the center of a door that is 27 1/2 inches wide, you will need to place the bar about 9 inches from each edge; this estimate can be used as a check on the | Algebraic expressions Algebraic equations Numerical expressions Numerical equations | 7.EE.3 Solve multi-step real-life and mathematical problems posed with positive and negative rational numbers in any form. Convert if necessary. | Positive number Negative number Rational number Equation Expression Decimal Percent Fraction |
| EE | Solve real-life and mathematical problems using numerical and algebraic expressions and equations. | exact computation. 7.EE.3 Solve multi-step real-life and mathematical problems posed with positive and negative rational numbers in any form (whole numbers, fractions, and decimals), using tools strategically. Apply properties of operations to calculate with numbers in any form; convert between forms as appropriate; and assess the reasonableness of answers using mental computation and estimation strategies. For example: If a woman making \$25 an hour gets a 10% raise, she will make an additional 1/10 of her salary an hour, or \$2.50, for a new salary of \$27.50. If you want to place a towel bar 9 3/4 inches long in the center of a door that is 27 1/2 inches wide, you will need to place the bar about 9 inches from each edge; this estimate can be used as a check on the exact computation. | | 7.EE.3 Evaluate the reasonableness of answers using mental computation and estimation strategies. Convert if necessary. | Estimation Resonable expression equation decimal percent fraction |

| Domain | Cluster | Common Core Standard | Content | Skills | Academic Vocabulary |
|--------|--|--|---|--|--|
| EE | mathematical problems using | 7.EE.4 Use variables to represent quantities in a real- world or mathematical problem, and construct simple equations and inequalities to solve problems by reasoning about the quantities. | Algebraic expressions and algebraic inequalities | 7.EE.4 Choose variables to represent quantities in a real-world or mathematical problem and construct simple equations and inequalities to solve problems by reasoning about the quantities. | Rational numbers Variables Equation Inequality |
| EE | mathematical problems using numerical and algebraic expressions and equations. | 7.EE.4a Solve word problems leading to equations of the form $px + q = r$ and $p(x + q) = r$, where p , q , and r are specific rational numbers. Solve equations of these forms fluently. Compare an algebraic solution to an arithmetic solution, identifying the sequence of the operations used in each approach. For example, the perimeter of a rectangle is 54 cm. Its length is 6 cm. What is its width? | Algebraic and numerical equations | | Rational numbers Variables Equation Inequality |
| EE | mathematical problems using numerical and algebraic expressions and equations. | 7.EE.4a Solve word problems leading to equations of the form $px + q = r$ and $p(x + q) = r$, where p , q , and r are specific rational numbers. Solve equations of these forms fluently. Compare an algebraic solution to an arithmetic solution, identifying the sequence of the operations used in each approach. For example, the perimeter of a rectangle is 54 cm. Its length is 6 cm. What is its width? | Algebraic and numerical equations | 7.EE.4a Compare an algebraic solution to an arithmetic solution by Identifying the sequence of the operations used in each approach. | Rational numbers Variables Equation Inequality |
| EE | mathematical problems using numerical and algebraic expressions and equations. | 7.EE.4b Solve word problems leading to inequalities of the form $px + q > r$ or $px + q < r$, where p , q , and r are specific rational numbers. Graph the solution set of the inequality and interpret it in the context of the problem. For example: As a salesperson, you are paid \$50 per week plus \$3 per sale. This week you want your pay to be at least \$100. Write an inequality for the number of sales you need to make, and describe the solutions. | Algebraic inequalities Numerical inequalities | | Inequality Rational numbers Solution set Infinite |

| Domain | Cluster | Common Core Standard | Content | Skills | Academic Vocabulary |
|--------|---|---|--|--------------------|---|
| EE | Solve real-life and mathematical problems using numerical and algebraic expressions and equations. | 7.EE.4b Solve word problems leading to inequalities of the form px + q > r or px + q < r, where p, q, and r are specific rational numbers. Graph the solution set of the inequality and interpret it in the context of the problem. For example: As a salesperson, you are paid \$50 per week plus \$3 per sale. This week you want your pay to be at least \$100. Write an inequality for the number of sales you need to make, and describe the solutions. | Algebraic inequalities Numerical inequalities | | Solution set Inequality Number line Infinite |
| EE | Solve real-life and mathematical problems using numerical and algebraic expressions and equations. | 7.EE.4b Solve word problems leading to inequalities of the form px + q > r or px + q < r, where p, q, and r are specific rational numbers. Graph the solution set of the inequality and interpret it in the context of the problem. For example: As a salesperson, you are paid \$50 per week plus \$3 per sale. This week you want your pay to be at least \$100. Write an inequality for the number of sales you need to make, and describe the solutions. | Numerical inequalities | | Solution set Inequality Number line Infinite |
| G | Draw construct, and describe geometrical figures and describe the relationships between them. | 7.G.1 Solve problems involving scale drawings of geometric figures, including computing actual lengths and areas from a scale drawing and reproducing a scale drawing at a different scale. | Proportional Relationships | | Scale drawing Scale factor Scale model Scale bar Scale distance |
| G | Draw construct, and describe geometrical figures and describe the relationships between them. | 7.G.1 Solve problems involving scale drawings of geometric figures, including computing actual lengths and areas from a scale drawing and reproducing a scale drawing at a different scale. | Proportional Relationships | geometric figures. | Scale drawing Scale factor Scale model Scale bar Scale distance |

| Domain | Cluster | Common Core Standard | Content | Skills | Academic Vocabulary |
|--------|----------------------|---|-----------------|---|------------------------|
| G | Draw construct, and | 7.G.2 Draw (freehand, with ruler and protractor, and with | Two dimensional | 7.G.2 Draw (freehand, with a ruler and protractor, and | Unique triangle |
| | describe geometrical | technology) geometric shapes with given conditions. | shapes | with technology) geometric shapes with given | Side |
| | figures and describe | Focus on constructing triangles from three measures of | | conditions. | Lengths |
| | the relationships | angles or sides, noticing when the conditions determine a | | | Angle measure |
| | between them. | unique triangle, more than one triangle, or no triangle. | | | Conjecture |
| | | | | | Conditions |
| G | Draw construct, and | 7.G.2 Draw (freehand, with ruler and protractor, and with | Two dimensional | 7.G.2 Determine if measures of three angles or sides | Unique triangle |
| | describe geometrical | technology) geometric shapes with given conditions. | shapes | create a unique triangle, more than one triangle, or no | Side |
| | figures and describe | Focus on constructing triangles from three measures of | | triangle. | Lengths |
| | the relationships | angles or sides, noticing when the conditions determine a | | | Angle measure |
| | between them. | unique triangle, more than one triangle, or no triangle. | | | Conjecture |
| | | | | | Conditions |
| G | Draw construct, and | 7.G.2 Draw (freehand, with ruler and protractor, and with | Two dimensional | 7.G.2 Construct triangles from three measures of | Unique triangle |
| | describe geometrical | technology) geometric shapes with given conditions. | shapes | angles or sides. | Side |
| | figures and describe | Focus on constructing triangles from three measures of | | | Lengths |
| | the relationships | angles or sides, noticing when the conditions determine a | | | Angle measure |
| | between them. | unique triangle, more than one triangle, or no triangle. | | | Conjecture |
| | | | | | Conditions |

| Domain | Cluster | Common Core Standard | Content | Skills | Academic Vocabulary |
|--------|----------------------|--|-----------------|---|--------------------------|
| G | Draw construct, and | 7.G.3 Describe the two-dimensional figures that result | Two dimensional | 7.G.3 Identify and describe the two-dimensional | Two dimensional shapes |
| | describe geometrical | from slicing three-dimensional figures, as in plane | shapes | figures that result from slicing three-dimensional | Three dimensional shapes |
| | figures and describe | sections of right rectangular prisms and right rectangular | | figures | Coplanar |
| | the relationships | pyramids. | | | Parallel |
| | between them. | | | | Solid |
| | | | | | Polyhedron |
| | | | | | Diagonal |
| | | | | | Slicing |
| | | | | | Cross section |
| | | | | | Vertical slice |
| | | | | | Angled slice |
| | | | | | Horizontal cross section |
| | | | | | Base |
| | | | | | Face |
| | | | | | Edges |
| | | | | | Vertices |
| | | | | | Rectangular Prisms |
| | | | | | Rectangular Pryamid |
| G | Solve real-life and | 7.G.4 Know the formulas for the area and circumference | Two dimensional | 7.G.4 Express and apply the formulas for the area and | Circumference |
| | mathematical | of a circle and use them to solve problems; give an | shapes | circumference of a circle. | Area |
| | problems involving | informal derivation of the relationship between the | | | Pi |
| | angle measure, area, | circumference and area of a circle. | | | Diameter |
| | surface area, and | | | | Radius |
| | volume. | | | | |
| | Solve real-life and | 7.G.4 Know the formulas for the area and circumference | Two dimensional | 7.G.4 Informally derive the relationship between the | Circumference |
| | mathematical | of a circle and use them to solve problems; give an | shapes | circumference and area of a circle. | Area |
| | problems involving | informal derivation of the relationship between the | | | Pi |
| | angle measure, area, | circumference and area of a circle. | | | Diameter |
| | surface area, and | | | | Radius |
| | volume. | | | | |

| Domain | Cluster | Common Core Standard | Content | Skills | Academic Vocabulary |
|--------|---|---|--------------|---|--|
| G | mathematical problems involving | 7.G.5 Use facts about supplementary, complementary, vertical, and adjacent angles in a multi-step problem to write and solve simple equations for an unknown angle in a figure. | | 7.G.5 Write and solve simple equations for an unknown angle in a figure using facts about supplementary, complementary, vertical, and adjacent angles in a multi-step problem. | Equations Supplementary angles Complementary angles Vertical angles Adjacent angles |
| | mathematical problems involving | 7.G.6 Solve real-world and mathematical problems involving area, volume and surface area of two- and three-dimensional objects composed of triangles, quadrilaterals, polygons, cubes, and right prisms. | Area | 7.G.6 Solve real-world and mathematical problems involving area of two-dimensional objects composed of triangles, quadrilaterals, and polygons. | Compose figures Triangles Quadrilaterals Polygons Area Square unit |
| | mathematical problems involving | 7.G.6 Solve real-world and mathematical problems involving area, volume and surface area of two- and three-dimensional objects composed of triangles, quadrilaterals, polygons, cubes, and right prisms. | Volume | 7.G.6 Solve real-world and mathematical problems involving volume of three-dimensional objects composed of cubes and right prisms. | Three dimensional shapes Cubes units Right prism Volume Square units |
| | mathematical problems involving | 7.G.6 Solve real-world and mathematical problems involving area, volume and surface area of two- and three-dimensional objects composed of triangles, quadrilaterals, polygons, cubes, and right prisms. | Surface Area | 7.G.6 Solve real-world and mathematical problems involving surface area of three-dimensional objects composed of cubes and right prisms. | Three dimensional shapes Cubes units Right prism Volume Surface area Square units |
| SP | to draw inferences about a population. | 7.SP.1 Understand that statistics can be used to gain information about a population by examining a sample of the population; generalizations about a population from a sample are valid only if the sample is representative of that population. Understand that random sampling tends to produce representative samples and support valid inferences. | | 7.SP.1 Explain that statistics can be used to gain information about a population by examining a representative sample of the population. | Data Survey Statistic Sample Population Representative sample Inferences Visual overlap |

| Domain | Cluster | Common Core Standard | Content | Skills | Academic Vocabulary |
|--------|--|---|-----------------|--|---|
| SP | | 7.SP.1 Understand that statistics can be used to gain information about a population by examining a sample of the population; generalizations about a population from a sample are valid only if the sample is representative of that population. Understand that random sampling tends to produce representative samples and support valid inferences. | | from a sample are valid only if the sample is representative of that population. | Data Generalization Survey Statistic Sample Population Representative sample Valid Inferences |
| SP | Use random sampling to draw inferences about a population. | 7.SP.1 Understand that statistics can be used to gain information about a population by examining a sample of the population; generalizations about a population from a sample are valid only if the sample is representative of that population. Understand that random sampling tends to produce representative samples and support valid inferences. | | | Data Generalization Survey Statistic Sample Population Representative sample Valid Inferences Random sampling |
| SP | Use random sampling to draw inferences about a population. | 7.SP.2 Use data from a random sample to draw inferences about a population with an unknown characteristic of interest. Generate multiple samples (or simulated samples) of the same size to gauge the variation in estimates or predictions. For example, estimate the mean word length in a book by randomly sampling words from the book; predict the winner of a school election based on randomly sampled survey data. Gauge how far off the estimate or prediction might be. | Random Sampling | characteristic of interest using data from a random sample. | Unknown characteristic of interest Statistic Population Random Sample Representative Sample |

| Domain | Cluster | Common Core Standard | Content | Skills | Academic Vocabulary |
|--------|--|---|-----------------|---|---|
| SP | Use random sampling to draw inferences about a population. | 7.SP.2 Use data from a random sample to draw inferences about a population with an unknown characteristic of interest. Generate multiple samples (or simulated samples) of the same size to gauge the variation in estimates or predictions. For example, estimate the mean word length in a book by randomly sampling words from the book; predict the winner of a school election based on randomly sampled survey data. Gauge how far off the estimate or prediction might be. | Random Sampling | samples) of the same size to gauge the variation in | Simulated sample Variation Predication |
| SP | Draw informal comparative inferences about two populations. | 7.SP.3 Informally assess the degree of visual overlap of two numerical data distributions with similar variabilities, measuring the difference between the centers by expressing it as a multiple of a measure of variability. For example, the mean height of players on the basketball team is 10 cm greater than the mean height of players on the soccer team, about twice the variability (mean absolute deviation) on either team; on a dot plot, the separation between the two distributions of heights is noticeable. | | | Dot plot Box-and-whisker plot Interquartile Range Mean absolute variation Visual overlap Numerical data distribution Similar variablities |
| SP | Draw informal comparative inferences about two populations. | 7.SP.3 Informally assess the degree of visual overlap of two numerical data distributions with similar variabilities, measuring the difference between the centers by expressing it as a multiple of a measure of variability. For example, the mean height of players on the basketball team is 10 cm greater than the mean height of players on the soccer team, about twice the variability (mean absolute deviation) on either team; on a dot plot, the separation between the two distributions of heights is noticeable. | | 7.SP.3 Measure the difference between the centers of two numerical data distributions with similar variabilities by expressing it as a multiple of a measure of variability. | Visual overlap Numerical data distribution Similar variablities Measure of variability(mean absolute variation), dot plot |

| Domain | Cluster | Common Core Standard | Content | Skills | Academic Vocabulary |
|--------|---|--|---------------------------------|---|---|
| SP | Draw informal comparative inferences about two populations. | 7.SP.4 Use measures of center and measures of variability for numerical data from random samples to draw informal comparative inferences about two populations. For example, decide whether the words in a chapter of a seventh-grade science book are generally longer than the words in a chapter of a fourth-grade science book. | Statistics and data analysis | | Population Random sample Measures of center (mean, median, mode) Range Measures of variability for Numerical data Comparative inferences |
| SP | Investigate chance processes and develop, use, and evaluate probability models. | 7.SP.5 Understand that the probability of a chance event is a number between 0 and 1 that expresses the likelihood of the event occurring. Larger numbers indicate greater likelihood. A probability near 0 indicates an unlikely event, a probability around 1/2 indicates an event that is neither unlikely nor likely, and a probability near 1 indicates a likely event. | Probability | 7.SP.5 Explain that the probability of a chance event is a number between 0 and 1 that expresses the likelihood of the event occurring. Larger numbers indicate greater likelihood. A probability near 0 indicates an unlikely event, a probability around 1/2 indicates an event that is neither unlikely nor likely, and a probability near 1 indicates a likely event. | Probability Simple events Outcome Likely event Unlikely event Certainty Impossible |
| SP | Investigate chance processes and develop, use, and evaluate probability models. | · | Probability | 7.SP.6 Approximate the probability of a chance event by collecting data on the chance process that produces it and observing its long-run relative frequency, and predict the approximate relative frequency given the probability. | Probability |
| SP | Investigate chance processes and develop, use, and evaluate probability models. | | Probability | 7.SP.7 | |

| Domain | Cluster | Common Core Standard | Content | Skills | Academic Vocabulary |
|--------|---|---|-------------|--|---|
| SP | Investigate chance processes and develop, use, and evaluate probability models. | 7.SP.7a Develop a uniform probability model by assigning equal probability to all outcomes, and use the model to determine probabilities of events. For example, if a student is selected at random from a class, find the probability that Jane will be selected and the probability that a girl will be selected. | Probability | assigning equal probability to all outcomes, and use the model to determine probabilities of events. | Probability Outcomes Probability model Equally likely Theoretical probability |
| SP | Investigate chance processes and develop, use, and evaluate probability models. | 7.SP.7b Develop a probability model (which may not be uniform) by observing frequencies in data generated from a chance process. For example, find the approximate probability that a spinning penny will land heads up or that a tossed paper cup will land open-end down. Do the outcomes for the spinning penny appear to be equally likely based on the observed frequencies? | Probability | | Probability model Observed frequencies Experimental probability Discrepancy |
| SP | Investigate chance processes and develop, use, and evaluate probability models. | 7.SP.8 Find probabilities of compound events using organized lists, tables, tree diagrams, and simulation. | Probability | 7.SP.8 | |
| SP | Investigate chance processes and develop, use, and evaluate probability models. | 7.SP.8a Understand that, just as with simple events, the probability of a compound event is the fraction of outcomes in the sample space for which the compound event occurs. | Probability | probability of a compound event is the fraction of outcomes in the sample space for which the compound event occurs. | Probability Simple events Outcomes Compound events Sample space |
| SP | Investigate chance processes and develop, use, and evaluate probability models. | 7.SP.8b Represent sample spaces for compound events using methods such as organized lists, tables and tree diagrams. For an event described in everyday language (e.g., "rolling double sixes"), identify the outcomes in the sample space which compose the event. | Probability | 7.SP.8b Represent sample spaces for compound events using methods such as organized lists, tables and tree diagrams. | |

| Domain | Cluster | Common Core Standard | Content | Skills | Academic Vocabulary |
|--------|----------------------|---|-------------|---|------------------------|
| SP | Investigate chance | 7.SP.8c Design and use a simulation to generate | Probability | 7.SP.8c Design and apply a simulation to generate | Probability |
| | processes and | frequencies for compound events. For example, use | | frequencies for compound events. | Simulation |
| | develop, use, and | random digits as a simulation tool to approximate the | | | Outcomes |
| | evaluate probability | answer to the question: If 40% of donors have type A | | | Compount events |
| | models. | blood, what is the probability that it will take at least 4 | | | Frequencies |
| | | donors to find one with type A blood? | | | |